arXiv:1509.07644v2 [math.NT] 10 Sep 2016

SHIFTED CONVOLUTION SUMS OF GL_3 CUSP FORMS WITH θ-SERIES

QINGFENG SUN

ABSTRACT. Let $A_f(1,n)$ be the normalized Fourier coefficients of a Hecke-Maass cusp form f for $SL_3(\mathbb{Z})$ and

$$r_3(n) = \#\left\{(n_1,n_2,n_3)\in\mathbb{Z}^3 : n_1^2+n_2^2+n_3^2=n\right\}.$$

Let $1\le h\le X$ and $\phi(x)$ be a smooth function compactly supported on $[1/2,1]$. It is proved that for any $\varepsilon>0$,

$$\sum_{n\geq 1} A_f(1,n+h)r_3(n)\phi\left(\frac{n}{X}\right) \ll_{f,\varepsilon} X^{\frac{3}{2}-\frac{1}{8}+\varepsilon}$$

uniformly with respect to the shift h.

CONTENTS

1. Introduction 1
2. Preliminaries on GL_3 cusp forms 5
3. Proof of Theorem 1.1 7
4. Proof of Proposition 3.2 17
5. Estimation of the character sum $\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)$ 18

References 22

1. INTRODUCTION

While the shifted convolution problems involving GL_2 automorphic forms are investigated intensively and deep results and applications have been established (see for example [2], [3], [7], [10], [14], [15], [18], [21], [25], [29]), the situation becomes much harder for GL_3 automorphic forms. For the cube of the Riemann zeta-function $\zeta^3(s)$, whose coefficients are $\tau_3(n)$, where $\tau_\ell(n)=\sum_{n_1n_2\cdots n_\ell=n}1$, Pitt [27], [28] first proved that, for $r>0$ an integer,

$$\sum_{n\leq X}\tau_3(n)a_g(rn-1) \ll_{g,\varepsilon} r^{\frac{5}{24}}X^{\frac{71}{72}+\varepsilon}, \tag{1.1}$$

where $a_g(n)$ are the normalized Fourier coefficients of a holomorphic cusp form for $SL_2(\mathbb{Z})$. Recently, Munshi [26] improved (1.1) by showing that $r^{\frac{5}{24}}X^{\frac{71}{72}+\varepsilon}$ can be replaced by

---

Date: September 13, 2016.

Key words and phrases. Shifted convolution sum, GL_3 cusp form, θ-series.

$r^{\frac{2}{7}}X^{\frac{34}{35}+\varepsilon}$. His method is Jutila's version of the circle method combined with the idea of factorable moduli. Moreover, let $A_f(1,n)$ and $a_g(n)$ be the normalized Fourier coefficients of a Hecke-Maass cusp form for $SL_3(\mathbb{Z})$ and $SL_2(\mathbb{Z})$, respectively. Let $0 \leq h \leq X$ and $\varphi(x)$ be a smooth function compactly supported in $[1,2]$. He also proved by the same method that (see [25])

$$\sum_{n\geq 1} A_f(1,n)a_g(n+h)\varphi\left(\frac{n}{X}\right) \ll_{f,g,\varepsilon} X^{1-\frac{1}{20}+\varepsilon} \tag{1.2}$$

holds uniformly respect to the shift h.

Let

$$r_\ell(n) = \#\left\{(n_1,n_2,\dots,n_\ell)\in\mathbb{Z}^\ell : n_1^2+\cdots+n_\ell^2 = n\right\}.$$

Then $r_\ell(n)$ is the n-th Fourier coefficient of the modular form $\theta^\ell(z)$, where $\theta(z)$ is the classical Jacobi theta series

$$\theta(z) = \sum_{n\in\mathbb{Z}} e(n^2 z).$$

The shifted convolution sum

$$\mathcal{S}_h(X) = \sum_{n\leq X} a_g(n+h)r_\ell(n),$$

with $a_g(n)$ the normalized Fourier coefficients of a holomorphic cusp form of weight κ for $\Gamma_0(N)$, was first studied by Luo [19]. Precisely, Luo first established a Voronoi formula for $r_\ell(n)$ and then applied Poincaré series reduction and his Voronoi formula to prove that for $\ell \geq 2$ and $\kappa \geq \frac{\ell}{2}+3$,

$$\mathcal{S}_h(X) \ll_{h,g,\ell,\varepsilon} X^{\frac{\ell}{2}-\vartheta_\ell+\varepsilon}, \tag{1.3}$$

where

$$\vartheta_\ell = \frac{\ell-1}{4(g+1)}, \qquad g = \begin{cases} \frac{1+\ell}{2}, & \text{if } \ell \text{ odd}, \\ 1+\frac{\ell}{2}, & \text{if } \ell \text{ even}. \end{cases}$$

In particular, $\vartheta_2 = \frac{1}{12}$, $\vartheta_3 = \frac{1}{6}$. Recently, Lü, Wu and Zhai [22] improved Luo's result by the circle method and showed that (1.3) holds uniformly for $0 \leq h \leq X$ and all κ, and ϑ_ℓ can be taken as $\theta_3 = \frac{1}{4}$, $\vartheta_\ell = \frac{1}{2}\,(\ell \geq 4)$. Moreover, for $N=1$, they proved that $\theta_2 = \frac{1}{6}$, $\vartheta_\ell = \frac{2}{3}\,(\ell \geq 6)$.

Usually the smaller ℓ's are more interesting but the related shifted convolution problems are more difficult. In this paper, we want to prove the following result.

Theorem 1.1. *Let $A_f(1,n)$ be the normalized Fourier coefficients of a Hecke-Maass cusp form for $SL_3(\mathbb{Z})$ and $1 \leq h \leq X$. Suppose that $\phi(x)$ is a smooth function supported on $[1/2,1]$ satisfying $\phi^{(j)}(x) \ll_j 1$. Then for $X > 1$ and any $\varepsilon > 0$, we have*

$$\sum_{n\geq 1} A_f(1,n+h)r_3(n)\phi\left(\frac{n}{X}\right) \ll_{f,\varepsilon} X^{\frac{3}{2}-\frac{1}{8}+\varepsilon}$$

uniformly respect to the shift h.

Remark 1. Under the Ramanujan conjecture, we have $A_f(1,n) \ll_\varepsilon n^\varepsilon$. Then by the asymptotic formula in the three-dimensional sphere problem (see [9])

$$\sum_{n\le X} r_3(n) = \frac{4\pi}{3} X^{\frac{3}{2}} + O\left(X^{\frac{21}{32}}\right),$$

we have the (conditional) bound

$$\sum_{n\ge 1} A_f(1,n+h) r_3(n) \phi\left(\frac{n}{X}\right) \ll_\varepsilon X^{\frac{3}{2}+\varepsilon}. \tag{1.4}$$

Further, by the argument in [22] (see Section 3), we have

$$\sum_{n\ge 1} A_f(1,n+h) r_3(n) \phi\left(\frac{n}{X}\right) \ll_\varepsilon X^{\frac{3}{2}-\left(\frac{3}{4}-\vartheta\right)+\varepsilon},$$

where ϑ is the exponent in the estimate

$$\sum_{n\ge 1} A_f(1,n) e(n\alpha) \phi\left(\frac{n}{X}\right) \ll_{f,\varepsilon} X^{\vartheta+\varepsilon}$$

which is uniform in $\alpha \in \mathbb{R}$. Using the best result due to Miller [23], we can take $\vartheta = \frac{3}{4}$ and also obtain the same bound as in (1.4). The bound in Theorem 1.1 is unconditional and better than (1.4) by a power of $X^{\frac{1}{8}-\varepsilon}$.

Remark 2. Although the shifted convolution sum in question should be comparable to the sum in (1.2) in view of the Voronoi formula for $r_\ell(n)$ established recently by Luo [19], the methods in Munshi [25] which obtains nontrivial power saving from the structural advantage of Jutila's variation of the circle method, are not applicable in our situation. This is because that there is no good upper bound for the exponential twisted sum

$$\sum_{n\le X} r_\ell(n) e(n\alpha), \qquad \alpha \in \mathbb{R}$$

which is, however, necessary for Jutila's version of the circle method.

Remark 3. For $h = 0$, the bound in Theorem 1.1 still holds. Moreover, Theorem 1.1 can be generalized to the cases $r_\ell(n)$, $\ell \ge 4$ without any difficulty. It is also worth noting that replacing $A_f(1,n)$ by the triple divisor function $\tau_3(n)$, we can get a better exponent saving than $\frac{1}{8} - \varepsilon$ due to the bound $\tau_3(n) \ll n^\varepsilon$ for any $\varepsilon > 0$. In fact, we have

$$\sum_{n\ge 1} \tau_3(n+h) r_3(n) \phi\left(\frac{n}{X}\right) = 4\mathcal{C}_2\mathcal{I}_0 X^{\frac{3}{2}} + 4\mathcal{C}_1\mathcal{I}_1 X^{\frac{3}{2}} + 2\mathcal{C}_0\mathcal{I}_2 X^{\frac{3}{2}} + O_\varepsilon\left(X^{\frac{3}{2}-\frac{1}{4}+\varepsilon}\right),$$

where for $\ell = 0, 1, 2$,

$$\mathcal{C}_\ell := \mathcal{C}_\ell(h) = \sum_{q=1}^{\infty} \frac{1}{q^5} \sum_{n\mid q} n\tau(n) P_\ell(n,q) \sum_{\substack{a=1\\(a,q)=1}}^{q} e\left(\frac{ah}{q}\right) G(a,0;q)^3 S\left(-\overline{a},0;\frac{q}{n}\right),$$

and

$$\mathcal{I}_\ell := \mathcal{I}_\ell(X,h,\phi) = \int_{-\infty}^{\infty}\left(\int_0^\infty \phi\left(\frac{u}{X}\right)e(-\beta u)(\log(u+h))^\ell \mathrm{d}u\right)\left(\int_0^1 e(\beta X v^2)\mathrm{d}v\right)^3 \mathrm{d}\beta.$$

Here $\tau(n)=\sum_{d|n}1$ is the divisor function, $\overline{a}$ denotes the multiplicative inverse of a mod q, $S(a,b;c)$ is the classical Kloosterman sum, $G(a,b;q)$ is the Gauss sum

$$G(a,b;q)=\sum_{d \bmod q} e\left(\frac{ad^2+bd}{q}\right),$$

$P_0(n,q)=1$ and $P_\ell(n,q)$ ($\ell=1,2$) are given by

$$\begin{aligned}
P_1(n,q) &= \frac{5}{3}\log n - 3\log q + 3\gamma - \frac{1}{3\tau(n)}\sum_{d|n}\log d,\\
P_2(n,q) &= (\log n)^2 - 5\log q\log n + \frac{9}{2}(\log q)^2 + 3\gamma^2 - 3\gamma_1 + 7\gamma\log n - 9\gamma\log q\\
&\qquad + \frac{1}{\tau(n)}\left((\log n+\log q-5\gamma)\sum_{d|n}\log d - \frac{3}{2}\sum_{d|n}(\log d)^2\right)
\end{aligned}$$

with $\gamma := \lim_{s\to 1}\left(\zeta(s)-\frac{1}{s-1}\right)$ being the Euler constant and $\gamma_1 := -\frac{\mathrm{d}}{\mathrm{d}s}\left(\zeta(s)-\frac{1}{s-1}\right)\big|_{s=1}$ being the Stieltjes constant. The proof is very similar as that of Theorem 1.1 (see also [30]).

Let us briefly outline the method of the proof. Applying the classical Hardy-Littlewood-Kloosterman circle method to the shifted convolution sum in Theorem 1.1, we arrive at an expression which is convenient to apply the Voronoi formula for GL_3 and an asymptotic formula for the sum $\sum_{|n|\le X} e(\alpha n^2)$, $\alpha\in[0,1]$. After that we are left with eight sums (in principle, of same difficulty) involving higher-dimensional character sums. In fact, we need to consider the twisted exponential sum (see Section 5)

$$\widetilde{\mathscr{T}}(p)=\sum_{x\in\mathbb{F}_p^\times}\left(\frac{x}{p}\right)e\left(\frac{r_1hx-\overline{4}(4v+b_1^2+b_2^2+b_3^2)\overline{x}}{p}\right)S(-r_2\overline{x},r_3n;p),$$

where p is an odd prime, $(r_i,p)=1$, $i=1,2,3$, and $b_1,b_2,b_3,v,n\in\mathbb{Z}$. This type of character sums have been studied in the work of Adolphson and Sperber [1]. The main saving in Theorem 1.1 comes from the nontrivial estimation of these character sums.

Notation. Throughout the paper, the letters h, q, m and n, with or without subscript, denote integers. The letter ε is an arbitrarily small positive constant, not necessarily the same at different occurrences. The symbol $\ll_{a,b,c}$ denotes that the implied constant depends at most on a, b and c.

2. PRELIMINARIES ON GL_3 CUSP FORMS

Let f be a Hecke-Maass cusp form of type $\nu = (\nu_1, \nu_2)$ for $SL_3(\mathbb{Z})$, normalized so that the first Fourier coefficient is 1. Then f has a Fourier-Whittaker expansion (see [5])

$$f(z) = \sum_{\gamma \in U_2(\mathbb{Z})\backslash SL_2(\mathbb{Z})} \sum_{n_1=1}^{\infty} \sum_{n_2 \neq 0} \frac{A_f(n_1,n_2)}{n_1|n_2|} W_J\left(M\begin{pmatrix} \gamma & \\ & 1 \end{pmatrix} z, \nu, \psi_{1,1}\right)$$

where $U_2(\mathbb{Z})$ is the group of 2×2 upper triangular matrices with integer entries and ones on the diagonal, $W_J(z,\nu,\psi_{1,1})$ is the Jacquet-Whittaker function and $M = \mathrm{diag}(n_1|n_2|, n_1, 1)$. By Rankin-Selberg theory, the Fourier coefficients $A_f(n_1,n_2)$ satisfy

$$\sum_{n_2 \leq N} |A_f(n_1,n_2)| \ll_f N|n_1|, \qquad \sum_{n_1 \leq N} |A_f(n_1,n_2)| \ll_f N|n_2|. \tag{2.1}$$

Let

$$\mu_1 = -\nu_1 - 2\nu_2 + 1, \qquad \mu_2 = -\nu_1 + \nu_2, \qquad \mu_3 = 2\nu_1 + \nu_2 - 1.$$

The generalized Ramanujan conjecture asserts that $\mathrm{Re}(\mu_j) = 0$, $1 \leq j \leq 3$, while the current record bound due to Luo, Rudnick and Sarnak [20] is

$$|\mathrm{Re}(\mu_j)| \leq \frac{1}{2} - \frac{1}{10}, \quad 1 \leq j \leq 3. \tag{2.2}$$

Let $\phi(x)$ be a smooth function compactly supported on $(0,\infty)$ and denote by $\widetilde{\phi}(s)$ the Mellin transform of $\phi(x)$. For $k = 0, 1$, we define

$$\Phi_k(x) := \int_{\mathrm{Re}(s)=\sigma} (\pi^3 x)^{-s} \prod_{j=1}^{3} \frac{\Gamma\left(\frac{1+s+\mu_j+2k}{2}\right)}{\Gamma\left(\frac{-s-\mu_j}{2}\right)} \widetilde{\phi}(-s-k)\mathrm{d}s \tag{2.3}$$

with $\sigma > \max\limits_{1\leq j\leq 3}\{-1-\mathrm{Re}(\mu_j)-2k\}$. Set

$$\Phi^{\pm}(x) = \Phi_0(x) \pm \frac{1}{i\pi^3 x}\Phi_1(x). \tag{2.4}$$

Then we have the following Voronoi-type formula (see [6], [24]).

Lemma 2.1. *Let $A_f(m,n)$ be the Fourier coefficients of a Maass cusp form for $SL_3(\mathbb{Z})$. Suppose that $\phi(x) \in C_c^{\infty}(0,\infty)$. Let $a, q \in \mathbb{Z}$ with $q \geq 1$, $(a,q) = 1$ and $a\overline{a} \equiv 1(\mathrm{mod} q)$. Then*

$$\begin{aligned}\sum_{n\geq 1} A_f(m,n) e\left(\frac{an}{q}\right)\phi(n) &= \frac{q\pi^{-\frac{5}{2}}}{4i} \sum_{n_1|qm} \sum_{n_2=1}^{\infty} \frac{A_f(n_2,n_1)}{n_1 n_2} S(m\overline{a}, n_2; mqn_1^{-1})\Phi^{+}\left(\frac{n_1^2 n_2}{q^3 m}\right) \\ &+ \frac{q\pi^{-\frac{5}{2}}}{4i} \sum_{n_1|qm} \sum_{n_2=1}^{\infty} \frac{A_f(n_1,n_2)}{n_1 n_2} S(m\overline{a}, -n_2; mqn_1^{-1})\Phi^{-}\left(\frac{n_1^2 n_2}{q^3 m}\right),\end{aligned}$$

where $S(m,n;c)$ is the classical Kloosterman sum.

The functions $\Phi^{\pm}(x)$ has the following properties.

Lemma 2.2. *Suppose that $\phi(x)$ is a smooth function of compact support in $[AX, BX]$, where $X > 0$ and $B > A > 0$, satisfying $\phi^{(j)}(x) \ll_{A,B,j} P^j$ for any integer $j \geq 0$. Then for $x > 0$ and any integer $N \geq 0$, we have*

$$\Phi^{\pm}(x) \ll_{f,A,B,N,\varepsilon} (xX)^{-\varepsilon}(PX)^3 \left(\frac{x}{P^3X^2}\right)^{-N}.$$

Proof. Let $\Phi_k(x)$ be as in (2.3). Changing variable $s + k \to s$, we have, for $\sigma > \max_{1\leq j\leq 3}\{-1 - \mathrm{Re}(\mu_j) - k\}$,

$$\Phi_k(x) = (\pi^3 x)^k \int\limits_{\mathrm{Re}(s)=\sigma} (\pi^3 x)^{-s} \prod_{j=1}^{3} \frac{\Gamma\left(\frac{1+s+\mu_j+k}{2}\right)}{\Gamma\left(\frac{-s-\mu_j+k}{2}\right)} \widetilde{\phi}(-s)\mathrm{d}s, \tag{2.5}$$

where

$$\widetilde{\phi}(s) = \int_0^{\infty} \phi(u)u^{s-1}\mathrm{d}u.$$

Let $s = \sigma + it$. By partial integration j times, we have, for $\sigma \notin \mathbb{Z}$,

$$\widetilde{\phi}(-s) = \frac{1}{s(s-1)\cdots(s-j+1)} \int_0^{\infty} \phi^{(j)}(u)u^{-s+j-1}\mathrm{d}u \ll_{A,B,\sigma,j} X^{-\sigma}(PX)^j(1+|t|)^{-j}. \tag{2.6}$$

Moreover, by Stirling's formula,

$$\prod_{j=1}^{3} \frac{\Gamma\left(\frac{1+s+\mu_j+k}{2}\right)}{\Gamma\left(\frac{-s-\mu_j+k}{2}\right)} \ll_{f,\sigma} \prod_{j=1}^{3}(1+|t|)^{\sigma+\frac{1}{2}+\mathrm{Re}(\mu_j)} \ll_{f,\sigma} (1+|t|)^{3\sigma+\frac{3}{2}} \tag{2.7}$$

in view of the fact that $\sum_{j=1}^{3} \mu_j = 0$.

Moving the contour to $\sigma = N+\varepsilon$ with $N \in \mathbb{Z}^+ \cup 0$ and taking $j = 3N+3$, by (2.5)-(2.7), we have

$$\begin{aligned}
\Phi_k(x) &\ll_{f,A,B,\sigma,j} x^k(xX)^{-\sigma}(PX)^j \int_{-\infty}^{+\infty} (1+|t|)^{-j+3\sigma+\frac{3}{2}}\mathrm{d}t \\
&\ll_{f,A,B,N,\varepsilon} x^k(xX)^{-N-\varepsilon}(PX)^{3N+3} \int_{-\infty}^{+\infty} (1+|t|)^{-\frac{3}{2}+3\varepsilon}\mathrm{d}t \\
&\ll_{f,A,B,N,\varepsilon} x^k(xX)^{-\varepsilon}(PX)^3 \left(\frac{x}{P^3X^2}\right)^{-N}. \qquad (2.8)
\end{aligned}$$

Then Lemma 2.2 follows from (2.4) and (2.8). □

By Lemma 2.2, for any fixed $\varepsilon > 0$ and $xX \geq X^{\varepsilon}(PX)^3$, $\Phi^{\pm}(x)$ is negligibly small. For $xX \ll X^{\varepsilon}$, as in [17], we move the line of integration in (2.6) to $\sigma = -\frac{11}{20}$ (guaranteed by

(2.2)), by (2.6) with $j = 1$ and (2.7), we have

$$\Phi_k(x) \ll_{f,A,B} x^k (xX)^{\frac{11}{20}} (PX) \int_{-\infty}^{+\infty} (1+|t|)^{-\frac{23}{20}} \mathrm{d}t \ll_{f,A,B,\varepsilon} x^k P X^{1+\varepsilon},$$

and by (2.4),

$$\Phi^{\pm}(x) \ll_{f,A,B,\varepsilon} P X^{1+\varepsilon}. \tag{2.9}$$

For $xX \gg X^{\varepsilon}$, we shall use the following result (see [16], [32]). For $\mu_1 = \mu_2 = \mu_3 = 0$, this was proved in [12].

Lemma 2.3. *Suppose that $\phi(x)$ is a smooth function of compact support on $[AX, BX]$, where $X > 0$ and $B > A > 0$. Then for $x > 0$, $xX \gg 1$, $\ell \geq 2$ and $k = 0, 1$, we have*

$$\begin{aligned}\Phi_k(x) &= (\pi^3 x)^{k+1} \sum_{j=1}^{\ell} \int_0^{\infty} \phi(u) \left(a_k(j) e\left(3(xu)^{\frac{1}{3}}\right) + b_k(j) e\left(-3(xu)^{\frac{1}{3}}\right)\right) \frac{\mathrm{d}u}{(\pi^3 xu)^{\frac{j}{3}}} \\ &\quad + O_{f,A,B,\varepsilon,\ell}\left((\pi^3 x)^k (\pi^3 xX)^{-\frac{\ell}{3}+\frac{1}{2}+\varepsilon}\right),\end{aligned}$$

where $a_k(j)$, $b_k(j)$ are constants with

$$a_0(1) = -\frac{2\sqrt{3\pi}}{3}, \quad b_0(1) = \frac{2\sqrt{3\pi}}{3}, \quad a_0(1) = b_1(1) = -\frac{2\sqrt{3\pi}}{3} i.$$

3. PROOF OF THEOREM 1.1

Denote

$$\mathscr{S}_h(X) = \sum_{n \geq 1} A_f(1, n+h) r_3(n) \phi\left(\frac{n}{X}\right).$$

We first transform $\mathscr{S}_h(X)$ by the Hardy-Littlewood-Kloosterman circle method (see for example, [13], Section 11.4). Let

$$\mathscr{F}(\alpha) = \sum_{|m| \leq \sqrt{X}} e(\alpha m^2)$$

and

$$\mathscr{G}(\alpha) = \sum_{n \geq 1} A_f(1, n+h) e(-\alpha n) \phi\left(\frac{n}{X}\right). \tag{3.1}$$

Then $\mathscr{S}_h(X)$ can be written as

$$\mathscr{S}_h(X) = \int_0^1 \mathscr{F}^3(\alpha) \mathscr{G}(\alpha) \mathrm{d}\alpha.$$

Note that $\mathscr{F}^3(\alpha)\mathscr{G}(\alpha)$ is a periodic function of period 1. We have

$$\mathscr{S}_h(X) = \int_{-1/(Q+1)}^{1-1/(Q+1)} \mathscr{F}^3(\alpha)\mathscr{G}(\alpha)\mathrm{d}\alpha,$$

where $Q = [5\sqrt{X}]$. Dissecting the unit interval with Farey's points of order Q, we have

$$\mathscr{S}_h(X) = \sum_{q\le Q} \sum_{a=1}^{q}{}^{*} \int\limits_{\mathscr{M}(a,q)} \mathscr{F}^3\left(\frac{a}{q}+\beta\right)\mathscr{G}\left(\frac{a}{q}+\beta\right)\mathrm{d}\beta,$$

where the $*$ denotes the condition $(a,q)=1$,

$$\mathscr{M}(a,q) = \left[-\frac{1}{q(q+q')}, \frac{1}{q(q+q'')}\right],$$

$\frac{a'}{q'}$, $\frac{a}{q}$ and $\frac{a''}{q''}$ are consecutive Farey fractions and they are determined by the conditions

$$Q < q+q', q+q'' \le q+Q, \quad aq' \equiv 1(\mathrm{mod} q), \quad aq'' \equiv -1(\mathrm{mod} q).$$

Exchanging the order of the summation over a and the integration over β as in Heath-Brown [8], we have

$$\mathscr{S}_h(X) = \sum_{q\le Q} \int\limits_{|\beta|\le\frac{1}{qQ}} \sum_{v \bmod q} \varrho(v,q,\beta) \sum_{a=1}^{q}{}^{*} e\left(-\frac{\overline{a}v}{q}\right)\mathscr{F}^3\left(\frac{a}{q}+\beta\right)\mathscr{G}\left(\frac{a}{q}+\beta\right)\mathrm{d}\beta, \tag{3.2}$$

where $\varrho(v,q,\beta)$ satisfies

$$\varrho(v,q,\beta) \ll \frac{1}{1+|v|}. \tag{3.3}$$

For an asymptotic formula of $\mathscr{F}\left(\frac{a}{q}+\beta\right)$, we quote the following result (see Theorem 4.1 in [31] or Lemma 4.1 in [33]).

Lemma 3.1. *Suppose that* $(a,q)=1$, $q \le Q$ *and* $|\beta| \le 1/(qQ)$. *We have*

$$\mathscr{F}\left(\frac{a}{q}+\beta\right) = \frac{2G(a,0;q)}{q}\Psi_0(\beta) + \sum_{-\frac{3q}{2}<b\le\frac{3q}{2}} G(a,b;q)\Psi(b,q,\beta), \tag{3.4}$$

where $G(a,b;q)$ *is the Gauss sum*

$$G(a,b;q) = \sum_{x \bmod q} e\left(\frac{ax^2+bx}{q}\right), \tag{3.5}$$

$\Psi_0(\beta)$ *is the integral*

$$\Psi_0(\beta) = \int_0^{\sqrt{X}} e(\beta x^2)\mathrm{d}x, \tag{3.6}$$

and $\Psi(b,q,\beta)$ *satisfies*

$$\sum_{-\frac{3q}{2}<b\leq\frac{3q}{2}} |\Psi(b,q,\beta)| \ll \log(q+2). \tag{3.7}$$

For $\mathscr{G}(\alpha)$ in (3.1), we apply Lemma 2.1 with $\phi_\beta(x) = \phi\left(\frac{x-h}{X}\right)e(-\beta x)$ getting

$$\begin{aligned}\mathscr{G}\left(\frac{a}{q}+\beta\right) &= e\left(\frac{ha}{q}+h\beta\right)\sum_{n\geq 1} A_f(1,n)e\left(-\frac{an}{q}\right)\phi\left(\frac{n-h}{X}\right)e(-\beta n)\\ &= \frac{q\pi^{-\frac{5}{2}}}{4i}e\left(\frac{ha}{q}+h\beta\right)\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_2,n_1)}{n_1n_2}S\left(-\overline{a},n_2;\frac{q}{n_1}\right)\Phi_\beta^{+}\left(\frac{n_1^2n_2}{q^3}\right)\\ &\quad+\frac{q\pi^{-\frac{5}{2}}}{4i}e\left(\frac{ha}{q}+h\beta\right)\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_1,n_2)}{n_1n_2}S\left(-\overline{a},-n_2;\frac{q}{n_1}\right)\Phi_\beta^{-}\left(\frac{n_1^2n_2}{q^3}\right),\end{aligned} \tag{3.8}$$

where

$$\Phi_\beta^{\pm}(x) = \Phi_0(x,\beta) \pm \frac{1}{i\pi^3x}\Phi_1(x,\beta) \tag{3.9}$$

with

$$\Phi_k(x,\beta) = \int\limits_{\mathrm{Re}(s)=\sigma}(\pi^3x)^{-s}\prod_{j=1}^{3}\frac{\Gamma\left(\frac{1+s+\mu_j+2k}{2}\right)}{\Gamma\left(\frac{-s-\mu_j}{2}\right)}\widetilde{\phi_\beta}(-s-k)\mathrm{d}s. \tag{3.10}$$

By (3.4) and (3.8), we have

$$\sum_{a=1}^{q}{}^{*}\, e\left(-\frac{\overline{a}v}{q}\right)\mathscr{F}^3\left(\frac{a}{q}+\beta\right)\mathscr{G}\left(\frac{a}{q}+\beta\right) = \sum_{j=1}^{8}\mathscr{D}_j(v,q,\beta), \tag{3.11}$$

where

$$\begin{aligned}
\mathscr{D}_1(v,q,\beta) &= \frac{2}{\pi^{\frac{5}{2}}i}\frac{e(h\beta)\Psi_0^3(\beta)}{q^2}\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_2,n_1)}{n_1n_2}\Phi_\beta^{+}\left(\frac{n_1^2n_2}{q^3}\right)\mathscr{C}(0,0,0,n_1,n_2,h,v;q), \qquad (3.12)\\
\mathscr{D}_2(v,q,\beta) &= \frac{2}{\pi^{\frac{5}{2}}i}\frac{e(h\beta)\Psi_0^3(\beta)}{q^2}\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_1,n_2)}{n_1n_2}\Phi_\beta^{-}\left(\frac{n_1^2n_2}{q^3}\right)\\
&\quad\times\mathscr{C}(0,0,0,n_1,-n_2,h,v;q),\\
\mathscr{D}_3(v,q,\beta) &= \frac{3}{\pi^{\frac{5}{2}}i}\frac{e(h\beta)\Psi_0^2(\beta)}{q}\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_2,n_1)}{n_1n_2}\Phi_\beta^{+}\left(\frac{n_1^2n_2}{q^3}\right)\\
&\quad\times\sum_{-\frac{3q}{2}<b\le\frac{3q}{2}}\Psi(b,q,\beta)\mathscr{C}(0,0,b,n_1,n_2,h,v;q), \qquad (3.13)\\
\mathscr{D}_4(v,q,\beta) &= \frac{3}{\pi^{\frac{5}{2}}i}\frac{e(h\beta)\Psi_0^2(\beta)}{q}\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_1,n_2)}{n_1n_2}\Phi_\beta^{-}\left(\frac{n_1^2n_2}{q^3}\right)\\
&\quad\times\sum_{-\frac{3q}{2}<b\le\frac{3q}{2}}\Psi(b,q,\beta)\mathscr{C}(0,0,b,n_1,-n_2,h,v;q),\\
\mathscr{D}_5(v,q,\beta) &= \frac{3}{2\pi^{\frac{5}{2}}i}e(h\beta)\Psi_0(\beta)\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_2,n_1)}{n_1n_2}\Phi_\beta^{+}\left(\frac{n_1^2n_2}{q^3}\right)\sum_{\substack{-\frac{3q}{2}<b_j\le\frac{3q}{2}\\ j=1,2}}\Psi(b_1,q,\beta)\Psi(b_2,q,\beta)\\
&\quad\times\mathscr{C}(0,b_1,b_2,n_1,n_2,h,v;q), \qquad (3.14)\\
\mathscr{D}_6(v,q,\beta) &= \frac{3}{2\pi^{\frac{5}{2}}i}e(h\beta)\Psi_0(\beta)\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_1,n_2)}{n_1n_2}\Phi_\beta^{-}\left(\frac{n_1^2n_2}{q^3}\right)\sum_{\substack{-\frac{3q}{2}<b_j\le\frac{3q}{2}\\ j=1,2}}\Psi(b_1,q,\beta)\Psi(b_2,q,\beta)\\
&\quad\times\mathscr{C}(0,b_1,b_2,n_1,-n_2,h,v;q),
\end{aligned}$$

$$\begin{aligned}
\mathscr{D}_7(v,q,\beta) &= \frac{q}{4\pi^{\frac{5}{2}}i}e(h\beta)\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_2,n_1)}{n_1n_2}\Phi_\beta^{+}\left(\frac{n_1^2n_2}{q^3}\right)\sum_{\substack{-\frac{3q}{2}<b_j\le\frac{3q}{2}\\ 1\le j\le 3}}\Psi(b_1,q,\beta)\Psi(b_2,q,\beta)\Psi(b_3,q,\beta)\\
&\quad\times\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q), \qquad (3.15)\\
\mathscr{D}_8(v,q,\beta) &= \frac{q}{4\pi^{\frac{5}{2}}i}e(h\beta)\sum_{n_1|q}\sum_{n_2=1}^{\infty}\frac{A_f(n_1,n_2)}{n_1n_2}\Phi_\beta^{-}\left(\frac{n_1^2n_2}{q^3}\right)\sum_{\substack{-\frac{3q}{2}<b_j\le\frac{3q}{2}\\ 1\le j\le 3}}\Psi(b_1,q,\beta)\Psi(b_2,q,\beta)\Psi(b_3,q,\beta)\\
&\quad\times\mathscr{C}(b_1,b_2,b_3,n_1,-n_2,h,v;q)
\end{aligned}$$

with

$$\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)=\sideset{}{^*}\sum_{a=1}^{q} e\left(\frac{ah-\overline{a}v}{q}\right)G(a,b_1;q)G(a,b_2;q)G(a,b_3;q)S\left(-\overline{a},n_2;\frac{q}{n_1}\right). \tag{3.16}$$

We only estimate the contributions from $\mathscr{D}_1$, $\mathscr{D}_3$, $\mathscr{D}_5$, $\mathscr{D}_7$, and the contributions from $\mathscr{D}_2$, $\mathscr{D}_4$, $\mathscr{D}_6$, $\mathscr{D}_8$ can be estimated similarly.

The following propositions will be proved in the next section.

Proposition 3.2. *For any $\varepsilon>0$, we have*

$$\sum_{n_2=1}^{\infty}\frac{|A_f(n_2,n_1)|}{n_1n_2}\left|\Phi_\beta^{\pm}\left(\frac{n_1^2n_2}{q^3}\right)\right|\ll_{f,\varepsilon}X^{\varepsilon}(1+|\beta|X)^2 \tag{3.17}$$

and

$$\sum_{n_2=1}^{\infty}\frac{|A_f(n_1,n_2)|}{n_1n_2}\left|\Phi_\beta^{\pm}\left(\frac{n_1^2n_2}{q^3}\right)\right|\ll_{f,\varepsilon}X^{\varepsilon}(1+|\beta|X)^2. \tag{3.18}$$

Proposition 3.3. *Let q_1 be the largest factor of q such that $q_1|n_1$ and $(q_1,q/q_1)=1$. Let q_2 be the largest factor of q/q_1 such that $q_2|n_1^{\infty}$ and $\left(q_2,\frac{q}{q_1q_2}\right)=1$. Let $q=q_1q_2q_3'q_3''$, $(q_3',2q_3'')=1$, q_3' square-free and $4q_3''$ square-full. Then for any $\varepsilon>0$, we have*

$$\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)\ll_{\varepsilon}\frac{(q_1q_2q_3'')^{3+\varepsilon}q_3'^{\frac{5}{2}+\varepsilon}(h,q_3')^{\frac{1}{2}}}{\sqrt{n_1}}.$$

By the second derivative test and the trivial estimation, $\Psi_0(\beta)$ in (3.6) is bounded by

$$\Psi_0(\beta)\ll\left(\frac{X}{1+|\beta|X}\right)^{\frac{1}{2}}. \tag{3.19}$$

Note that the condition $\frac{n_1^2n_2}{q^3}X<X^{\varepsilon}(1+|\beta|X)^3$ with $|\beta|\le 1/(qQ)$ implies that $n_1^2n_2\ll X^{2+\varepsilon}/Q^3\ll X^{\frac{1}{2}+\varepsilon}$.

Let q be as in Proposition 3.3. Denote $q_0 = q_2 q_3''$. Note that q_0 is square-full. By (3.12), (3.17), (3.19), Lemma 2.2 and Proposition 3.3, we have

$$\begin{aligned}
\mathscr{D}_1(v,q,\beta) &\ll_{f,\varepsilon} \frac{1}{q^2}\left(\frac{X}{1+|\beta|X}\right)^{\frac{3}{2}} \sum_{n_1|q} \sum_{\frac{n_1^2 n_2}{q^3}X < X^{\varepsilon}(1+|\beta|X)^3} \frac{|A_f(n_2,n_1)|}{n_1 n_2}\left|\Phi_\beta^+\left(\frac{n_1^2 n_2}{q^3}\right)\right| \\
&\quad \times |\mathscr{C}(0,0,0,n_1,n_2,h,v;q)| + 1 \\
&\ll_{f,\varepsilon} X^{\varepsilon}\left(\frac{X}{1+|\beta|X}\right)^{\frac{3}{2}} \sum_{\substack{n_1 \ll X^{\frac{1}{4}+\varepsilon} \\ n_1 \equiv 0(\bmod q_1)}} \frac{q_1 q_0 q_3'^{\frac{1}{2}}(h,q_3')^{\frac{1}{2}}}{\sqrt{n_1}} \sum_{n_2=1}^{\infty} \frac{|A_f(n_2,n_1)|}{n_1 n_2}\left|\Phi_\beta^+\left(\frac{n_1^2 n_2}{q^3}\right)\right| + 1 \\
&\ll_{f,\varepsilon} X^{\frac{3}{2}+\varepsilon}(1+|\beta|X)^{\frac{1}{2}} \sum_{\substack{n_1 \ll X^{\frac{1}{4}+\varepsilon} \\ n_1 \equiv 0(\bmod q_1)}} \frac{q_1 q_0 q_3'^{\frac{1}{2}}(h,q_3')^{\frac{1}{2}}}{\sqrt{n_1}}.
\end{aligned}$$

By (3.3) and the estimate above, we have

$$\begin{aligned}
&\sum_{q\le Q}\int_{|\beta|\le\frac{1}{qQ}}\sum_{v \bmod q}\varrho(v,q,\beta)\mathscr{D}_1(v,q,\beta)\mathrm{d}\beta\\
\ll_{f,\varepsilon}\;& X^{\frac32+\varepsilon}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}\sum_{\substack{q_3'\le Q/q_1\\ q_3'\ \mathrm{square-free}}}\sum_{\substack{q_0\le Q/(q_1q_3')\\ 4q_0\ \mathrm{square-full}}} q_1q_0{q_3'}^{\frac12}(h,q_3')^{\frac12}\int_{|\beta|\le\frac{1}{q_1q_0q_3'Q}}(1+|\beta|X)^{\frac12}\mathrm{d}\beta\\
\ll_{f,\varepsilon}\;& X^{\frac32+\varepsilon}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}\sum_{\substack{q_3'\le Q/q_1\\ q_3'\ \mathrm{square-free}}}\sum_{\substack{q_0\le Q/(q_1q_3')\\ 4q_0\ \mathrm{square-full}}} q_1q_0{q_3'}^{\frac12}(h,q_3')^{\frac12}\left(\frac{1}{q_1q_0q_3'Q}+\frac{X^{\frac12}}{(q_1q_0q_3'Q)^{\frac32}}\right)\\
\ll_{f,\varepsilon}\;& \frac{X^{\frac32+\varepsilon}}{Q}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}\sum_{q_3'\le Q/q_1}(h,q_3')^{\frac12}{q_3'}^{-\frac12}\sum_{\substack{q_0\le Q/(q_1q_3')\\ 4q_0\ \mathrm{square-full}}}1\\
&+\frac{X^{2+\varepsilon}}{Q^{\frac32}}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}q_1^{-\frac12}\sum_{q_3'\le Q/q_1}(h,q_3')^{\frac12}{q_3'}^{-1}\sum_{\substack{q_0\le Q/(q_1q_3')\\ 4q_0\ \mathrm{square-full}}}q_0^{-\frac12}\\
\ll_{f,\varepsilon}\;& \frac{X^{\frac32+\varepsilon}}{Q}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}\sum_{q_3'\le Q/q_1}(h,q_3')^{\frac12}{q_3'}^{-\frac12}\left(\frac{Q}{q_1q_3'}\right)^{\frac12}\\
&+\frac{X^{2+\varepsilon}}{Q^{\frac32}}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}q_1^{-\frac12}\sum_{q_3'\le Q/q_1}(h,q_3')^{\frac12}{q_3'}^{-1}\\
\ll_{f,\varepsilon}\;& \frac{X^{\frac32+\varepsilon}}{Q^{\frac12}}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}q_1^{-\frac12}\sum_{d|h}d^{-\frac12}\sum_{q_4\le Q/q_1d}q_4^{-1}\\
&+\frac{X^{2+\varepsilon}}{Q^{\frac32}}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}q_1^{-\frac12}\sum_{d|h}d^{-\frac12}\sum_{q_4\le Q/(q_1d)}q_4^{-1}\\
\ll_{f,\varepsilon}\;& \frac{X^{\frac32+\frac18+\varepsilon}}{Q^{\frac12}}+\frac{X^{2+\frac18+\varepsilon}}{Q^{\frac32}}\\
\ll_{f,\varepsilon}\;& X^{\frac32-\frac18+\varepsilon} \qquad (3.20)
\end{aligned}$$

uniformly for $1\le h\le X$.

Further, by (3.7), (3.13), Lemma 2 and Proposition 3.3, we have

$$\begin{aligned}
\mathscr{D}_3(v,q,\beta) \ll_{f,\varepsilon}\; & \frac{1}{q}\frac{X}{1+|\beta|X}\sum_{n_1|q}\sum_{\frac{n_1^2n_2}{q^3}X<X^{\varepsilon}(1+|\beta|X)^3}\frac{|A_f(n_2,n_1)|}{n_1n_2}\left|\Phi_\beta^+\left(\frac{n_1^2n_2}{q^3}\right)\right| \\
& \times\sum_{-\frac{3q}{2}<b\leq\frac{3q}{2}}|\Psi(b,q,\beta)||\mathscr{C}(0,0,b,n_1,n_2,h,v;q)|+1 \\
\ll_{f,\varepsilon}\; & \frac{X^{1+\varepsilon}}{1+|\beta|X}\sum_{\substack{n_1\ll X^{\frac{1}{4}+\varepsilon}\\ n_1\equiv 0(\bmod q_1)}}\frac{q_1^2q_0^2{q_3'}^{\frac{3}{2}}(h,q_3')^{\frac{1}{2}}}{\sqrt{n_1}}\sum_{n_2=1}^{\infty}\frac{|A_f(n_2,n_1)|}{n_1n_2}\left|\Phi_\beta^+\left(\frac{n_1^2n_2}{q^3}\right)\right|+1 \\
\ll_{f,\varepsilon}\; & X^{1+\varepsilon}(1+|\beta|X)\sum_{\substack{n_1\ll X^{\frac{1}{4}+\varepsilon}\\ n_1\equiv 0(\bmod q_1)}}\frac{q_1^2q_0^2{q_3'}^{\frac{3}{2}}(h,q_3')^{\frac{1}{2}}}{\sqrt{n_1}}.
\end{aligned}$$

It follows from this estimate and (3.3) that

$$\begin{aligned}
& \sum_{q\leq Q}\int_{|\beta|\leq\frac{1}{qQ}}\sum_{v \bmod q}\varrho(v,q,\beta)\mathscr{D}_3(v,q,\beta)\mathrm{d}\beta \\
\ll_{f,\varepsilon}\; & X^{1+\varepsilon}\sum_{n_1\ll X^{\frac{1}{4}+\varepsilon}}n_1^{-\frac{1}{2}}\sum_{q_1|n_1}\sum_{\substack{q_3'\leq Q/q_1\\ q_3'\ \text{square-free}}}\sum_{\substack{q_0\leq Q/(q_1q_3')\\ 4q_0\ \text{square-full}}}q_1^2q_0^2{q_3'}^{\frac{3}{2}}(h,q_3')^{\frac{1}{2}}\int_{|\beta|\leq\frac{1}{q_1q_0q_3'Q}}(1+|\beta|X)\mathrm{d}\beta \\
\ll_{f,\varepsilon}\; & X^{1+\varepsilon}\sum_{n_1\ll X^{\frac{1}{4}+\varepsilon}}n_1^{-\frac{1}{2}}\sum_{q_1|n_1}\sum_{\substack{q_3'\leq Q/q_1\\ q_3'\ \text{square-free}}}\sum_{\substack{q_0\leq Q/(q_1q_3')\\ 4q_0\ \text{square-full}}}q_1^2q_0^2{q_3'}^{\frac{3}{2}}(h,q_3')^{\frac{1}{2}}\left(\frac{1}{q_1q_0q_3'Q}+\frac{X}{(q_1q_0q_3'Q)^2}\right) \\
\ll_{f,\varepsilon}\; & \frac{X^{1+\varepsilon}}{Q}\sum_{n_1\ll X^{\frac{1}{4}+\varepsilon}}n_1^{-\frac{1}{2}}\sum_{q_1|n_1}q_1\sum_{q_3'\leq Q/q_1}(h,q_3')^{\frac{1}{2}}{q_3'}^{\frac{1}{2}}\left(\frac{Q}{q_1q_3'}\right)^{\frac{3}{2}} \\
& +\frac{X^{2+\varepsilon}}{Q^2}\sum_{n_1\ll X^{\frac{1}{4}+\varepsilon}}n_1^{-\frac{1}{2}}\sum_{q_1|n_1}\sum_{q_3'\leq Q/q_1}(h,q_3')^{\frac{1}{2}}{q_3'}^{-\frac{1}{2}}\left(\frac{Q}{q_1q_3'}\right)^{\frac{1}{2}} \\
\ll_{f,\varepsilon}\; & X^{1+\frac{1}{8}+\varepsilon}Q^{\frac{1}{2}}+\frac{X^{2+\frac{1}{8}+\varepsilon}}{Q^{\frac{3}{2}}} \\
\ll_{f,\varepsilon}\; & X^{\frac{3}{2}-\frac{1}{8}+\varepsilon} \qquad (3.21)
\end{aligned}$$

uniformly for $1\leq h\leq X$.

Moreover, by (3.7), (3.14), Lemma 2.2 and Propositions 3.3,

$$\begin{aligned}
\mathscr{D}_5(v,q,\beta) \quad &\ll_{f,\varepsilon} \quad \left(\frac{X}{1+|\beta|X}\right)^{\frac{1}{2}} \sum_{n_1|q} \sum_{\frac{n_1^2 n_2}{q^3} X < X^{\varepsilon}(1+|\beta|X)^3} \frac{|A_f(n_2,n_1)|}{n_1 n_2} \left|\Phi_{\beta}^{+}\left(\frac{n_1^2 n_2}{q^3}\right)\right| \\
&\qquad \times \sum_{\substack{-\frac{3q}{2} < b_j \le \frac{3q}{2} \\ j=1,2}} |\Psi(b_1,q,\beta)|\Psi(b_2,q,\beta)||\mathscr{C}(0,b_1,b_2,n_1,n_2,h,v;q)| + 1 \\
&\ll_{f,\varepsilon} \quad X^{\varepsilon}\left(\frac{X}{1+|\beta|X}\right)^{\frac{1}{2}} \sum_{\substack{n_1 \ll X^{\frac{1}{4}+\varepsilon} \\ n_1 \equiv 0(\bmod q_1)}} n_1^{-\frac{1}{2}} q_1^3 q_0^3 q_3'^{\frac{5}{2}} (h,q_3')^{\frac{1}{2}} \sum_{n_2=1}^{\infty} \frac{|A_f(n_2,n_1)|}{n_1 n_2} \left|\Phi_{\beta}^{+}\left(\frac{n_1^2 n_2}{q^3}\right)\right| + 1 \\
&\ll_{f,\varepsilon} \quad X^{\frac{1}{2}+\varepsilon}(1+|\beta|X)^{\frac{3}{2}} \sum_{\substack{n_1 \ll X^{\frac{1}{4}+\varepsilon} \\ n_1 \equiv 0(\bmod q_1)}} n_1^{-\frac{1}{2}} q_1^3 q_0^3 q_3'^{\frac{5}{2}} (h,q_3')^{\frac{1}{2}}.
\end{aligned}$$

Applying (3.3) again, we obtain

$$\begin{aligned}
&\sum_{q \le Q} \int_{|\beta| \le \frac{1}{qQ}} \sum_{v \bmod q} \varrho(v,q,\beta) \mathscr{D}_5(v,q,\beta) \mathrm{d}\beta \\
\ll_{f,\varepsilon} \quad & X^{\frac{1}{2}+\varepsilon} \sum_{n_1 \ll X^{\frac{1}{4}+\varepsilon}} n_1^{-\frac{1}{2}} \sum_{q_1|n_1} \sum_{\substack{q_3' \le Q/q_1 \\ q_3' \text{ square-free}}} \sum_{\substack{q_0 \le Q/(q_1 q_3') \\ 4q_0 \text{ square-full}}} q_1^3 q_0^3 q_3'^{\frac{5}{2}} (h,q_3')^{\frac{1}{2}} \int_{|\beta| \le \frac{1}{q_1 q_0 q_3' Q}} (1+|\beta|X)^{\frac{3}{2}} \mathrm{d}\beta \\
\ll_{f,\varepsilon} \quad & X^{\frac{1}{2}+\varepsilon} \sum_{n_1 \ll X^{\frac{1}{4}+\varepsilon}} n_1^{-\frac{1}{2}} \sum_{q_1|n_1} \sum_{\substack{q_3' \le Q/q_1 \\ q_3' \text{ square-free}}} \sum_{\substack{q_0 \le Q/(q_1 q_3') \\ 4q_0 \text{ square-full}}} q_1^3 q_0^3 q_3'^{\frac{5}{2}} (h,q_3')^{\frac{1}{2}} \left(\frac{1}{q_1 q_0 q_3' Q} + \frac{X^{\frac{3}{2}}}{(q_1 q_0 q_3' Q)^{\frac{5}{2}}}\right) \\
\ll_{f,\varepsilon} \quad & \frac{X^{\frac{1}{2}+\varepsilon}}{Q} \sum_{n_1 \ll X^{\frac{1}{4}+\varepsilon}} n_1^{-\frac{1}{2}} \sum_{q_1|n_1} q_1^2 \sum_{q_3' \le Q/q_1} (h,q_3')^{\frac{1}{2}} q_3'^{\frac{3}{2}} \left(\frac{Q}{q_1 q_3'}\right)^{\frac{5}{2}} \\
& + \frac{X^{2+\varepsilon}}{Q^{\frac{5}{2}}} \sum_{n_1 \ll X^{\frac{1}{4}+\varepsilon}} n_1^{-\frac{1}{2}} \sum_{q_1|n_1} q_1^{\frac{1}{2}} \sum_{q_3' \le Q/q_1} (h,q_3')^{\frac{1}{2}} \frac{Q}{q_1 q_3'} \\
\ll_{f,\varepsilon} \quad & X^{\frac{1}{2}+\frac{1}{8}+\varepsilon} Q^{\frac{3}{2}} + \frac{X^{2+\frac{1}{8}+\varepsilon}}{Q^{\frac{3}{2}}} \\
\ll_{f,\varepsilon} \quad & X^{\frac{3}{2}-\frac{1}{8}+\varepsilon} \qquad (3.22)
\end{aligned}$$

uniformly for $1 \le h \le X$.

Lastly, by (3.7), (3.15), Lemma 2.2 and Propositions 3.3, we have

$$\begin{aligned}
\mathscr{D}_7(v,q,\beta) \ &\ll_{f,\varepsilon} \ q\sum_{n_1|q}\sum_{\frac{n_1^2n_2}{q^3}X<X^{\varepsilon}(1+|\beta|X)^3}\frac{|A_f(n_2,n_1)|}{n_1n_2}\left|\Phi_\beta^+\left(\frac{n_1^2n_2}{q^3}\right)\right| \\
&\qquad \times\sum_{\substack{-\frac{3q}{2}<b_j\leq\frac{3q}{2}\\ 1\leq j\leq 3}}|\Psi(b_1,q,\beta)||\Psi(b_2,q,\beta)||\Psi(b_3,q,\beta)||\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)|+1 \\
&\ll_{f,\varepsilon} \ X^{\varepsilon}\sum_{\substack{n_1\ll X^{\frac14+\varepsilon}\\ n_1\equiv 0(\bmod q_1)}} n_1^{-\frac12}q_1^4q_0^4q_3'^{\frac72}(h,q_3')^{\frac12}\sum_{n_2=1}^{\infty}\frac{|A_f(n_2,n_1)|}{n_1n_2}\left|\Phi_\beta^+\left(\frac{n_1^2n_2}{q^3}\right)\right|+1 \\
&\ll_{f,\varepsilon} \ X^{\varepsilon}(1+|\beta|X)^2\sum_{\substack{n_1\ll X^{\frac14+\varepsilon}\\ n_1\equiv 0(\bmod q_1)}} n_1^{-\frac12}q_1^4q_0^4q_3'^{\frac72}(h,q_3')^{\frac12}.
\end{aligned}$$

It follows that

$$\begin{aligned}
&\sum_{q\leq Q}\int_{|\beta|\leq\frac{1}{qQ}}\sum_{v \bmod q}\varrho(v,q,\beta)\mathscr{D}_7(v,q,\beta)\mathrm{d}\beta \\
\ll_{f,\varepsilon} \ & X^{\varepsilon}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}\sum_{\substack{q_3'\leq Q/q_1\\ q_3' \text{ square-free}}}\sum_{\substack{q_0\leq Q/(q_1q_3')\\ 4q_0 \text{ square-full}}} q_1^4q_0^4q_3'^{\frac72}(h,q_3')^{\frac12}\int_{|\beta|\leq\frac{1}{q_1q_0q_3'Q}}(1+|\beta|X)^2\mathrm{d}\beta \\
\ll_{f,\varepsilon} \ & X^{\varepsilon}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}\sum_{\substack{q_3'\leq Q/q_1\\ q_3' \text{ square-free}}}\sum_{\substack{q_0\leq Q/(q_1q_3')\\ 4q_0 \text{ square-full}}} q_1^4q_0^4q_3'^{\frac72}(h,q_3')^{\frac12}\left(\frac{1}{q_1q_0q_3'Q}+\frac{X^2}{(q_1q_0q_3'Q)^3}\right) \\
\ll_{f,\varepsilon} \ & \frac{X^{\varepsilon}}{Q}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}q_1^3\sum_{q_3'\leq Q/q_1}(h,q_3')^{\frac12}q_3'^{\frac52}\left(\frac{Q}{q_1q_3'}\right)^{\frac72} \\
& +\frac{X^{2+\varepsilon}}{Q^3}\sum_{n_1\ll X^{\frac14+\varepsilon}} n_1^{-\frac12}\sum_{q_1|n_1}q_1\sum_{q_3'\leq Q/q_1}(h,q_3')^{\frac12}q_3'^{\frac12}\left(\frac{Q}{q_1q_3'}\right)^{\frac32} \\
\ll_{f,\varepsilon} \ & X^{\frac18+\varepsilon}Q^{\frac52}+\frac{X^{2+\frac18+\varepsilon}}{Q^{\frac32}} \\
\ll_{f,\varepsilon} \ & X^{\frac32-\frac18+\varepsilon} \qquad (3.23)
\end{aligned}$$

uniformly for $1\leq h\leq X$. By (3.2), (3.11) and (3.20)-(3.23), Theorem 1.1 follows.

4. PROOF OF PROPOSITION 3.2

We only prove (3.17) and (3.18) can be proved similarly. Recall $\Phi_\beta^\pm(x)$ in (3.9) which we relabel as

$$\Phi_\beta^\pm(x) = \Phi_0(x,\beta) \pm \frac{1}{i\pi^3 x}\Phi_1(x,\beta), \tag{4.1}$$

where for $\sigma > \max\limits_{1\le j\le 3}\{-1-\mathrm{Re}(\mu_j)-2k\}$,

$$\Phi_k(x,\beta) = \int\limits_{\mathrm{Re}(s)=\sigma} (\pi^3 x)^{-s} \prod_{j=1}^{3} \frac{\Gamma\left(\frac{1+s+\mu_j+2k}{2}\right)}{\Gamma\left(\frac{-s-\mu_j}{2}\right)} \widetilde{\phi_\beta}(-s-k)\mathrm{d}s \tag{4.2}$$

with $\phi_\beta(x) = \phi\left(\frac{x-h}{X}\right)e(-\beta x)$. Note that

$$\phi_\beta^{(j)}(x) \ll_j \left(\frac{1+|\beta|X}{X}\right)^j.$$

By Lemma 2.2, we have

$$\sum_{n_2=1}^{\infty} \frac{|A_f(n_2,n_1)|}{n_1 n_2}\left|\Phi_\beta^\pm\left(\frac{n_1^2 n_2}{q^3}\right)\right| = \sum_{\frac{n_1^2 n_2}{q^3}X < X^\varepsilon(1+|\beta|X)^3} \frac{|A_f(n_2,n_1)|}{n_1 n_2}\left|\Phi_\beta^\pm\left(\frac{n_1^2 n_2}{q^3}\right)\right| + O_{f,\varepsilon}(1) \tag{4.3}$$

By (2.1) and (2.9), we have

$$\begin{aligned} \sum_{\frac{n_1^2 n_2}{q^3}X \le X^\varepsilon} \frac{|A_f(n_2,n_1)|}{n_2}\left|\Phi_\beta^\pm\left(\frac{n_1^2 n_2}{q^3}\right)\right| &\ll_{f,\varepsilon} X^\varepsilon(1+|\beta|X) \max_{1\le T\le \frac{q^3X^\varepsilon}{n_1^2 X}} \frac{1}{T}\sum_{T\le n_2\le 2T}|A_f(n_2,n_1)| \\ &\ll_{f,\varepsilon} X^\varepsilon n_1(1+|\beta|X). \end{aligned} \tag{4.4}$$

For $xX > X^\varepsilon$, by Lemma 2.3, we have

$$\begin{aligned} \Phi_k(x,\beta) &= (\pi^3 x)^{k+1}\sum_{j=1}^{\ell}\int_0^\infty \phi\left(\frac{u-h}{X}\right)e(-\beta u)\left(a_k(j)e\left(3(xu)^{\frac{1}{3}}\right) + b_k(j)e\left(-3(xu)^{\frac{1}{3}}\right)\right)\frac{\mathrm{d}u}{(\pi^3 xu)^{\frac{j}{3}}} \\ &\quad + O_{f,\varepsilon,\ell}\left((\pi^3 x)^k(\pi^3 xX)^{-\frac{\ell}{3}+\frac{1}{2}+\varepsilon}\right), \end{aligned}$$

where $a_k(j)$, $b_k(j)$ are constants, and by (4.1),

$$\Phi_\beta^\pm(x) \ll_{f,\varepsilon,\ell} x\sum_{j=1}^{\ell} x^{-\frac{j}{3}}\left(|\mathcal{I}_j(x,\beta)| + |\mathcal{J}_j(x,\beta)|\right) + (xX)^{-\frac{\ell}{3}+\frac{1}{2}+\varepsilon}, \tag{4.5}$$

with

$$\begin{aligned}\mathcal{I}_j(x,\beta) &= \int_0^\infty u^{-\frac{j}{3}}\phi\left(\frac{u-h}{X}\right)e(-\beta u)e\left(3(xu)^{\frac{1}{3}}\right)\mathrm{d}u,\\ \mathcal{J}_j(x,\beta) &= \int_0^\infty u^{-\frac{j}{3}}\phi\left(\frac{u-h}{X}\right)e(-\beta u)e\left(-3(xu)^{\frac{1}{3}}\right)\mathrm{d}u.\end{aligned}$$

By partial integration twice, we have

$$\begin{aligned}\mathcal{I}_j(x,\beta) &= \frac{1}{\left(2\pi i x^{\frac{1}{3}}\right)^2}\int_0^\infty \left(u^{\frac{2}{3}}\left(u^{\frac{2-j}{3}}\phi\left(\frac{u-h}{X}\right)e(-\beta u)\right)'\right)' e\left(3(xu)^{\frac{1}{3}}\right)\mathrm{d}u\\ &\ll (xX)^{-\frac{2}{3}}X^{1-\frac{j}{3}}(1+|\beta|X)^2. \qquad (4.6)\end{aligned}$$

Similarly,

$$\begin{aligned}\mathcal{J}_j(x,\beta) &= \frac{1}{\left(2\pi i x^{\frac{1}{3}}\right)^2}\int_0^\infty \left(u^{\frac{2}{3}}\left(u^{\frac{2-j}{3}}\phi\left(\frac{u-h}{X}\right)e(-\beta u)\right)'\right)' e\left(-3(xu)^{\frac{1}{3}}\right)\mathrm{d}u\\ &\ll (xX)^{-\frac{2}{3}}X^{1-\frac{j}{3}}(1+|\beta|X)^2. \qquad (4.7)\end{aligned}$$

Taking $\ell = 3$. By (4.5)-(4.7), we have

$$\Phi_\beta^{\pm}(x) \ll_{f,\varepsilon,\ell} (xX)^{\frac{1}{3}}(1+|\beta|X)^2\sum_{j=1}^{\ell}(xX)^{-\frac{j}{3}} + (xX)^{-\frac{\ell}{3}+\frac{1}{2}+\varepsilon} \ll_{f,\varepsilon} (1+|\beta|X)^2.$$

This estimate combined with (2.1) yields that

$$\begin{aligned}&\sum_{X^\varepsilon<\frac{n_1^2n_2}{q^3}X<X^\varepsilon(1+|\beta|X)^3}\frac{|A_f(n_2,n_1)|}{n_2}\left|\Phi_\beta^{\pm}\left(\frac{n_1^2n_2}{q^3}\right)\right|\\ \ll_{f,\varepsilon}\ & (1+|\beta|X)^2(\log X)\max_{\frac{q^3X^\varepsilon}{n_1^2X}<T<\frac{q^3X^\varepsilon(1+|\beta|X)^3}{n_1^2X}}\frac{1}{T}\sum_{T\le n_2\le 2T}|A_f(n_2,n_1)|\\ \ll_{f,\varepsilon}\ & X^\varepsilon n_1(1+|\beta|X)^2. \qquad (4.8)\end{aligned}$$

Then Proposition 3.2 follows from (4.3), (4.4) and (4.8).

5. ESTIMATION OF THE CHARACTER SUM $\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)$

Let $b_1,b_2,b_3,n_1,n_2,v\in\mathbb{Z}$, $n_1|q$ and $1\le h\le X$. We recall $\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)$ in (3.16) which we relabel as

$$\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)=\sideset{}{^*}\sum_{a \bmod q} e\left(\frac{ah-\overline{a}v}{q}\right)G(a,b_1;q)G(a,b_2;q)G(a,b_3;q)S\left(-\overline{a},n_2;\frac{q}{n_1}\right), \qquad (5.1)$$

where $S(m,n;c)$ is the classical Kloosterman sum and $G(a,b;q)$ is the Gauss sum in (3.5)

$$G(a,b;q)=\sum_{x \bmod q} e\left(\frac{ax^2+bx}{q}\right).$$

In this section, we shall prove Proposition 3.3. We need the following results for $G(a,b;q)$ (see Lemma 5.4.5 in [11])

Lemma 5.1. *(1) If $(q_1,q_2)=1$, then $G(a,b;q_1q_2)=G(aq_2,b;q_1)G(aq_1,b;q_2)$.*

(2) For $(a,q)=1$, the sum $G(a,b;q)$ has absolute value $\sqrt{q}$ if q is odd, $\sqrt{2q}$ if q is even, $q=2r$ and $ar+b$ is even.

(3) For $(2a,q)=1$, we have

$$G(a,0;q)=\left(\frac{a}{q}\right)\epsilon_q\sqrt{q},$$

where $\epsilon_q=\begin{cases} 1, & \text{if } q\equiv 1(\mathrm{mod} 4),\\ i, & \text{if } q\equiv -1(\mathrm{mod} 4).\end{cases}$

(4) For q odd, we have

$$G(a,b;q)=e\left(-\frac{\bar{4}\bar{a}b^2}{q}\right)G(a,0;q).$$

In the estimate for $\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)$, the main case of interest is when q is square-free and $n_1=1$. Thus we first extract from q the largest part related to n_1, the contribution of which to $\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)$ will be trivially estimated (using Weil's bound for Kloosterman sums). More precisely, let q_1 be the largest factor of q such that $q_1|n_1$ and $(q_1,q/q_1)=1$. Let q_2 be the largest factor of q/q_1 such that $q_2|n_1^{\infty}$ and $\left(q_2,\frac{q}{q_1q_2}\right)=1$. Note that q_2 is square-full and $n_1|q_1q_2$. Denote temporarily $q'=q_1q_2$ and $\widehat{q}=\frac{q'}{n_1}$. Let $q_3=\frac{q}{q_1q_2}$. Then any reduced residue class $a \bmod q$ can be written as $a=a_1q_3+a_2q'$ with inverse $\overline{a}=\overline{a}_1q_3\overline{q}_3^2+\overline{a}_2q'\overline{q'}^2$, where $a_1 \bmod q'$ and $a_2 \bmod q_3$ are some reduced residue classes modulo q' and q_3, respectively. Then by Lemma 5.1 (1), we have

$$\begin{aligned}
&\mathscr{C}(b_1,b_2,b_3,n_1,n_2,h,v;q)\\
={}&\sideset{}{^*}\sum_{a_1 \bmod q'} e\left(\frac{a_1h-\overline{a}_1\overline{q}_3^2v}{q'}\right)G(a_1q_3^2,b_1;q')G(a_1q_3^2,b_2;q')G(a_1q_3^2,b_3;q')S(-\overline{a_1q_3},n_2\overline{q_3}^2;\widehat{q})\\
&\times\sideset{}{^*}\sum_{a_2 \bmod q_3} e\left(\frac{a_2h-\overline{a}_2\overline{q'}^2v}{q_3}\right)G(a_2q'^2,b_1;q_3)G(a_2q'^2,b_2;q_3)G(a_2q'^2,b_3;q_3)S(-\overline{a_2q'},n_2\overline{\widehat{q}}^2;q_3)\\
={}&\sideset{}{^*}\sum_{a_1 \bmod q'} e\left(\frac{a_1\overline{q}_3^2h-\overline{a_1}v}{q'}\right)G(a_1,b_1;q')G(a_1,b_2;q')G(a_1,b_3;q')S(-\overline{a_1}q_3,n_2\overline{q_3}^2;\widehat{q})\\
&\times\sideset{}{^*}\sum_{a_2 \bmod q_3} e\left(\frac{a_2\overline{q'}^2h-\overline{a}_2v}{q_3}\right)G(a_2,b_1;q_3)G(a_2,b_2;q_3)G(a_2,b_3;q_3)S(-\overline{a_2}q',n_2\overline{\widehat{q}}^2;q_3)\\
:={}&\mathscr{C}^*(b_1,b_2,b_3,n_1,n_2,h,v;q')\mathscr{C}^{**}(b_1,b_2,b_3,n_1,n_2,h,v;q_3) \qquad (5.2)
\end{aligned}$$

say.

By Lemma 5.1 (2), we have $G(a,b;q) \ll \sqrt{q}$. This estimate combined with Weil's bound for Kloosterman sum gives

$$\mathscr{C}^*(b_1,b_2,b_3,n_1,n_2,h,v;q') \ll q'^{\frac{5}{2}}\left(\overline{a_1}q_3,n_2\overline{q_3}^2,\frac{q'}{n_1}\right)^{\frac{1}{2}}\left(\frac{q'}{n_1}\right)^{\frac{1}{2}}\tau\left(\frac{q'}{n_1}\right) \ll \frac{q_1^3q_2^3\tau(q_1q_2)}{\sqrt{n_1}}. \qquad (5.3)$$

Next we extract the square-full part from the remaining part of q, that is to say, we further factor q_3 as $q_3=q_3'q_3''$ with $(q_3',2q_3'')=1$, q_3' square-free and $4q_3''$ square-full. Then

$$\mathscr{C}^{**}(b_1,b_2,b_3,n_1,n_2,h,v;q_3)=\mathscr{C}_1^{**}(b_1,b_2,b_3,n_1,n_2,h,v;q_3')\mathscr{C}_2^{**}(b_1,b_2,b_3,n_1,n_2,h,v;q_3'') \qquad (5.4)$$

where

$$\begin{aligned}
\mathscr{C}_1^{**}(b_1,b_2,b_3,n_1,n_2,h,v;q_3') &= \sideset{}{^*}\sum_{\gamma \bmod q_3'} e\left(\frac{\gamma\overline{q'}^2\overline{q_3''}^2h-\overline{\gamma}v}{q_3'}\right)G(\gamma,b_1;q_3')G(\gamma,b_2;q_3')\\
&\quad G(\gamma,b_3;q_3')S(-\overline{\gamma}q'q_3'',n_2\overline{\widehat{q}}^2\overline{q_3''}^2;q_3')\\
\mathscr{C}_2^{**}(b_1,b_2,b_3,n_1,n_2,h,v;q_3'') &= \sideset{}{^*}\sum_{\gamma \bmod q_3''} e\left(\frac{\gamma\overline{q'}^2\overline{q_3'}^2h-\overline{\gamma}v}{q_3''}\right)G(\gamma,b_1;q_3'')G(\gamma,b_2;q_3'')\\
&\quad G(\gamma,b_3;q_3'')S(-\overline{\gamma}q'q_3',n_2\overline{\widehat{q}}^2\overline{q_3'}^2;q_3'').
\end{aligned}$$

By Lemma 5.1 (2) and Weil's bound for Kloosterman sum, we have

$$\mathscr{C}_2^{**}(b_1,b_2,b_3,n_1,n_2,h,v;q_3'') \ll q''^3\tau(q_3''). \qquad (5.5)$$

To estimate $\mathscr{C}_1^{**} := \mathscr{C}_1^{**}(b_1, b_2, b_3, n_1, n_2, h, v; q_3')$, we factor q_3' as $q_3' = p_1 p_2 \cdots p_s$, p_i prime. Then

$$\mathscr{C}_1^{**} = \prod_{i=1}^{s} \mathscr{T}(b_1, b_2, b_3, \overline{q'}^2 \overline{q_3''}^2 \overline{p_i'}^2 h, q' q_3'' p_i', n_2 \overline{\widehat{q}}^2 \overline{q_3''}^2 \overline{p_i'}^2; p_i), \tag{5.6}$$

where $p_i' = q_3'/p_i$ and

$$\mathscr{T}(b_1, b_2, b_3, r_1 h, r_2, r_3 n_2; p) = \sideset{}{^*}\sum_{x \bmod p} e\left(\frac{r_1 h x - v\overline{x}}{p}\right) G(x, b_1; p) G(x, b_2; p) G(x, b_3; p) S(-r_2\overline{x}, r_3 n_2; p)$$

with $(p, 2) = 1$ and $(r_i, p) = 1$, i=1,2,3.

By Lemma 5.1 (3), (4), we have

$$G(x, b_j; p) = e\left(-\frac{\overline{4}\overline{x} b_j^2}{p}\right) G(x, 0; p) = e\left(-\frac{\overline{4}\overline{x} b_j^2}{p}\right) \left(\frac{x}{p}\right) \epsilon_p \sqrt{p}.$$

Hence

$$\begin{aligned} &\mathscr{T}(b_1, b_2, b_3, r_1 h, r_2, r_3 n_2; p) \\ = \;& \epsilon_p^3 p^{\frac{3}{2}} \sideset{}{^*}\sum_{x \bmod p} \left(\frac{x}{p}\right) e\left(\frac{r_1 h x - \overline{4}(4v + b_1^2 + b_2^2 + b_3^2)\overline{x}}{p}\right) S(-r_2\overline{x}, r_3 n_2; p). \end{aligned} \tag{5.7}$$

Denote

$$\widetilde{\mathscr{T}}(p) = \sideset{}{^*}\sum_{x \bmod p} \left(\frac{x}{p}\right) e\left(\frac{r_1 h x - \overline{4}(4v + b_1^2 + b_2^2 + b_3^2)\overline{x}}{p}\right) S(-r_2\overline{x}, r_3 n_2; p).$$

By (5.1)-(5.7), Proposition 3.3 follows from the following lemma.

Lemma 5.2. *We have*

$$\widetilde{\mathscr{T}}(p) \ll (h, p)^{\frac{1}{2}} p.$$

Proof. If $p | h$, then by Weil's bound for Kloosterman sum, we have

$$\widetilde{\mathscr{T}}(p) \ll p^{\frac{3}{2}}. \tag{5.8}$$

If $p \nmid h$, $p | n_2$, then $S(-r_2\overline{x}, r_3 n_2; p) = -1$ and

$$\widetilde{\mathscr{T}}(p) = -\sideset{}{^*}\sum_{x \bmod p} \left(\frac{x}{p}\right) e\left(\frac{r_1 h x - \overline{4}(4v + b_1^2 + b_2^2 + b_3^2)\overline{x}}{p}\right) \ll p^{\frac{1}{2}} \tag{5.9}$$

by the bound for Salié sum (see [13], Corollary 4.10).

If $p \nmid h$, $p \nmid n_2$, we open the Kloosterman sum to obtain

$$\widetilde{\mathscr{T}}(p) = \sum_{x, y \in \mathbb{F}_p^\times} \left(\frac{x}{p}\right) e\left(\frac{r_1 h x - \overline{4}(4v + b_1^2 + b_2^2 + b_3^2)\overline{x} - r_2\overline{x} y + r_3 n_2 \overline{y}}{p}\right).$$

The square-root cancellation for such twisted character sums was established in the more general case in [1] (see also [4]). To apply their result to the special character sum $\widetilde{\mathscr{T}}(p)$, we consider the Newton polyhedron $\Delta(f)$ of $f(x,y) = r_1hx - \overline{4}(4v + b_1^2 + b_2^2 + b_3^2)x^{-1} - r_2x^{-1}y + r_3n_2y^{-1} \in \mathbb{F}_p^\times[x, y, (xy)^{-1}]$. If $p \nmid 4v + b_1^2 + b_2^2 + b_3^2$, then $\Delta(f)$ is the quadrilateral in $\mathbb{R}^2$ with vertices $(1,0)$, $(-1,0)$, $(-1,1)$ and $(0,-1)$. If $p \mid 4v + b_1^2 + b_2^2 + b_3^2$, then $\Delta(f)$ is the triangle in $\mathbb{R}^2$ with vertices $(1,0)$, $(-1,1)$ and $(0,-1)$. Thus in both cases $\dim\Delta(f) = 2$ and $(0,0)$ is an interior point of $\Delta(f)$. Moreover, for $p \nmid 4v + b_1^2 + b_2^2 + b_3^2$, each of the following eight polynomials

$$\begin{aligned} f_\sigma(x,y) &= r_1hx, -\overline{4}(4v + b_1^2 + b_2^2 + b_3^2)x^{-1}, -r_2x^{-1}y, r_3n_2y^{-1}, r_1hx - r_2x^{-1}y, r_1hx + r_3n_2y^{-1}, \\ &\quad -\overline{4}(4v + b_1^2 + b_2^2 + b_3^2)x^{-1} - r_2x^{-1}y, -\overline{4}(4v + b_1^2 + b_2^2 + b_3^2)x^{-1} + r_3n_2y^{-1} \end{aligned}$$

and for $p \mid 4v + b_1^2 + b_2^2 + b_3^2$, each of the following six polynomials

$$f_\sigma(x,y) = r_1hx, -r_2x^{-1}y, r_3n_2y^{-1}, r_1hx - r_2x^{-1}y, r_1hx + r_3n_2y^{-1}, -r_2x^{-1}y + r_3n_2y^{-1}$$

corresponding to the faces of $\Delta(f)$ not containing $(0,0)$, the locus of

$$\frac{\partial f_\sigma}{\partial x} = \frac{\partial f_\sigma}{\partial y} = 0$$

is empty in $\left(\overline{F}_p^\times\right)^2$. Thus in both cases f is nondegenerate with respect to $\Delta(f)$. By [1] or [4], we have

$$\widetilde{\mathscr{T}}(p) \ll p. \tag{5.10}$$

Then Lemma 5.2 follows from (5.8)-(5.10). □

ACKNOWLEDGEMENTS. The author would like to express heartfelt thanks to the referees for their important and useful comments. This work is supported by the National Natural Science Foundation of China (Grant No. 11101239) and Young Scholars Program of Shandong University, Weihai (Grant No. 2015WHWLJH04).

REFERENCES

[1] A. Adolphson and S. Sperber *Twisted exponential sums and Newton polyhedra*, J. reine angew. Math 443 (1993), 151-177.

[2] V. Blomer and G. Harcos, *The spectral decomposition of shifted convolution sums*, Duke Math. J. 144 (2008), 321-339.

[3] W. Duke, J. Friedlander and H. Iwaniec, *Bounds for automorphic L-functions*, Invent. Math. 112 (1993), 1-8.

[4] L. Fu, *Weights of twisted exponential sums*, Math. Z. 262 (2009), 449-472.

[5] D. Goldfeld, *Automorphic Forms and L-functions for the group $GL(n,\mathbb{R})$*, with an appendix by Kevin A. Broughan, Cambridge Studies in Advanced Mathematics 99, Cambridge University Press, Cambridge, 2006.

[6] D. Goldfeld and X. Li, *Voronoi formulas on $GL(n)$*, Int. Math. Res. Not. 2006, Art. ID 86295, 25 pp.

[7] G. Harcos and P. Michel, *The subconvexity problem for RankinCSelberg L-functions and equidistribution of Heegner points. II*, Inventiones mathematicae 163 (3) (2006): 581-655.

[8] D. R. Heath-Brown, *Cubic forms in ten variables*, Proc. London Math. Soc. 3 (2) (1983), 225-257.

[9] D. R. Heath-Brown, *Lattice points in the sphere*, Number theory in progress 2 (1997), 883-892.

[10] R. Holowinsky, *A sieve method for shifted convolution sums*, Duke Math. J. 146 (2009), 401-448.

[11] M. N. Huxley, *Area, lattice points, and exponential sums*, Oxford University Press, 1996.

[12] Iv A. Ivić, *On the ternary additive divisor problem and the six moment of the zeta-function*, Sieve methods, exponential sums, and their applications in number theory (Cardiff, 1995), 205-243, London Math. Soc. Lecture Nore Ser., 237, Cambridge Univ. Press, Cambridge, 1997.

[13] H. Iwaniec, *Topics in classical automorphic forms*, American Mathematical Soc., 1997.

[14] M. Jutila, *The additive divisor problem and its analogs for Fourier coefficients of cusp forms. I, II*, Math. Z. 223 (1996), 435-461, 225 (1997), 625-637.

[15] E. Kowalski, P. Michel and J. Vanderkam. *Rankin-Selberg L-functions in the level aspect*, Duke Math. J. 114 (2002), 123-191.

[16] X. Li, *The central value of the Rankin-Selberg L-functions*, GAFA 18 (2009), 1660-1695.

[17] X. Li, *Bounds for $GL(3)\times GL(2)$ L-functions and $GL(3)$ L-functions*, Ann. of Math. (2) 173 (2011), 301-336.

[18] Y. K. Lau, J. Y. Liu and Y. B. Ye, *A new bound $k^{2/3+\varepsilon}$ for Rankin-Selberg L-functions for Hecke congruence subgroups*, Int. Math. Res. Papers, 2006, 35090.

[19] W. Luo, *Shifted convolution of cusp-forms with θ-series*, Abh. Math. Semin. Univ. Hambg. 81 (2011), 45-53.

[20] W. Luo, Z. Rudnick and P. Sarnak, *On the generalized Ramanujan conjecture for $GL(n)$*, Automorphic forms, automorphic representations, and arithmetic (Fort Worth, TX, 1996), 301-310, Proc. Sympos. Pure Math. 66, Part 2, Amer. Math. Soc., Providence, RI, 1999.

[21] W. Luo and P. Sarnak, *Mass equidistribution for Hecke eigenforms*, Comm. Pure Appl. Math. 56 (2003), 874-891.

[22] G. Lü, J. Wu and W. Zhai, *Shifted convolution of cusp-forms with θ-series*, Ramanujan J. (2015), DOI 10.1007/s11139-015-9678-8.

[23] S.D. Miller, *Cancellation in additively twisted sums on GL (n)*, American journal of mathematics, 128 (3) (2006), 699-729.

[24] S.D. Miller and W. Schmid, *Automorphic distributions, L-functions, and Voronoi summation for $GL(3)$*, Ann. of Math. (2) 164 (2006), no. 2, 423-488.

[25] R. Munshi, *Shifted convolution sums for $GL(3)\times GL(2)$*, Duke Math. J. 162 (2013), 2345-2362.

[26] R. Munshi, *Shifted convolution of divisor function d_3 and Ramanujan τ function*, in The legacy of Srinivasa Ramanujan, volume 20 of Ramanujan Math. Soc. Lect. Notes Ser., 251-260. Ramanujan Math. Soc., Mysore, 2013.

[27] N. J. E. Pitt, *On shifted convolution sums for $\zeta^3(s)$ with automorphic L-functions*, Duke Math. J. 77 (1995), 383-406.

[28] N. J. E. Pitt, *On an analogue of Titchmarsh's divisor problem for holomorphic cusp forms*, Journal of the American Mathematical Society, 26 (3) (2013), 735-776.

[29] P. Sarnak, *Estimates for Rankin-Selberg L-functions and quantum unique ergodicity*, J. Funct. Anal. 184 (2001), 419-453.

[30] Q. F. Sun and D. Y. Zhang, *Sums of the triple divisor function over values of a ternary quadratic form*, arXiv:1510.06170.

[31] R. C. Vaughan, *The Hardy-Littlewood method*, Cambridge University Press, 1997.

[32] X. M. Ren and Y. B. Ye, *Asymptotic Voronoi's summation formulas and their duality for $SL_3(\mathbb{Z})$*, Number theoryarithmetic in Shangri-La, 213C236, Ser. Number Theory Appl., 8, World Sci. Publ., Hackensack, NJ, 2013.

[33] L. L. Zhao, *The sum of divisors of a quadratic form*, Acta Arith. 163 (2014), no. 2, 161C177.

School of Mathematics and Statistics, Shandong University, Weihai, Weihai, Shandong 264209, China
E-mail address: `qfsun@sdu.edu.cn`